

## Rapid Ecological Integrity Assessments in Wetlands of Riparian Areas in Maryland: Coastal Plain Region

### Condensed Field Guidance, Procedures, and Scoring Tables (to be used with separate Data Sheet)

Project/Site Name: \_\_\_\_\_

Sampling Date: \_\_\_\_\_

#### GENERAL GUIDANCE

-This Ecological Integrity Assessment uses information collected in the field and from online sources/imagery. **Additional background and information can be found in the referenced sections of the “Field Manual for Rapid Ecological Integrity Assessments of Wetlands in Riparian Areas in Maryland: Coastal Plain” (Manual).**

-Refer to Section 2 in the Manual for field visit preparation and how to identify the wetland assessment area (AA) or areas on the project site. Each AA should be evaluated and scored separately.

-You will need to use online resources to prepare for the site visit, complete some of the data sheet, and to complete the Landscape Assessment. An Excel file can provide some autofill features when used with the wetland delineation Excel file (see Manual).

#### PROCESS

-**Review the metrics and example photos in the Manual. In the field, use the data sheet (Excel or pdf) and this document simultaneously to score and evaluate features.** For each metric, review the guidance in this document and carry out the procedures indicated to collect data. Record your data on the data sheet, using the check boxes to indicate features present and filling in other required information where needed. Use the data that you recorded on the data sheet and the scoring tables in this document to determine a score for each metric. The scores can be entered on this document, but they need also to be recorded on the data sheet or in the Excel file. Enter all scores on the Scoring Form and follow the Manual instructions to calculate the Final Score.

-Scoring can vary due to the conditions expected for different Key Wildlife Habitats (Section 4.3). **Be sure to use the sections of the tables that correspond to the Key Wildlife Habitat being evaluated.**

-**NOTE:** All of the characteristics described for a given score category may not be present. **Assign the score to the category with the majority of features present.**

#### LANDSCAPE ASSESSMENT (Section 3)

Watershed features can impact habitat quality for the organisms in the project area. Natural habitats provide the greatest benefit for wetland buffers, which play a critical role in the condition of the wetland relative to key abiotic and biotic factors. One Landscape Assessment is done for the entire project area and will apply to each AA in the project area. **Most of the landscape-level assessments will be done in the office** using mapped features and aerial imagery as described in the Manual. However, additional features noted in the field that are not visible on available imagery may affect the assessment. **In the field, as you are traveling to and assessing the AA, make note of the features described below to supplement the in-office assessment. Record these observations on the data sheet.** If access to the buffer area is limited, scoring will need to rely more on aerial imagery as described in the Manual.

Landscape Features	Assess out to this distance from the outer edges of the proposed stream restoration project area (all AA are included in project area):	Note these features on the data sheet for use with information from aerial imagery:
Buffer Perimeter	10m (33 feet)	Natural and altered habitats (see table below)
Buffer Condition	100m (330 feet)	Natural and altered habitats (see table below)
Aquatic Context	300m (1000 feet)	Small-scale wetlands, such as Springs or Vernal Pools, or streams that may not be evident from aerial imagery or are newly formed
Comparative Size	n/a- assessment occurs for each AA in the project area	Deviations from aerial imagery that could affect wetland size estimation; source(s) of size reduction of the AA such as roads, impoundment, development, etc.

Examples of Land Covers Included in Natural Buffers	Examples of Land Covers Excluded from Natural Buffers (Altered Habitats)
Natural plant communities; naturally vegetated rights-of-way; natural swales and ditches; natural open water features including rivers, streams, and ponds created by beaver activity; wetlands	Parking lots; commercial and private developments and structures; roads (all types); intensive agriculture; intensive plantations; orchards; vineyards; railroads; planted pastures; planted hayfields; animal pastures; lawns; sports fields; traditional golf courses; fallow farm fields; ditches; stormwater ponds; ponds formed by unnatural blockages; culverts

## SITE DESCRIPTION AND ENVIRONMENTAL INFORMATION (Sections 4.1 and 4.2)

Provide a detailed description of the assessment area on the data sheet, including landscape setting, vegetation type, evidence of human or natural disturbance, and characteristics of the stream and other nearby features. **Note Landscape Position, Water Source, and Hydrological Regime** for the AA. If there is more than one water source, rank as P (primary), S (secondary), and T (tertiary). The Hydrological Regime usually matches the mapped wetland designation (see Manual for definitions).

## ASSIGNMENT OF AA TO KEY WILDLIFE HABITAT (Section 4.3) and Vegetation Indicators

Use the key below to determine the Key Wildlife Habitat (KWH) for the AA. Also indicate the stream type and, if possible, the community type/plant association. See the Manual for photos and complete descriptions. Lists of typical species in each stratum by KWH and indicator species by KWH are listed on the next page. These species lists may assist with KWH selection and will be used in the KWH and Vegetation Composition metrics in Section 4.6.

**1a.** Wetlands bordering streams and rivers with overland, non-tidal flooding regimes (i.e., floodplains). Distinct alluvial landforms (e.g., backswamps, levees, terraces) and indicators present (e.g., scour marks, recent sediment deposition, vegetation damaged/bent in one direction, soils with alternating deposits, channel banks with flood marks). Likely to be 3<sup>rd</sup> order and higher but may be braided systems. Structurally and compositionally diverse vegetation present ranging from closed mixed forests to floodplain pools to open, beaver-created pools with floating aquatics.....**COASTAL PLAIN FLOODPLAIN** HGM Class: Riverine

**1b.** Wetlands primarily controlled via groundwater discharge often associated with depressional and slope geomorphic features as well as the margins of small stream (1<sup>st</sup> and 2<sup>nd</sup> order) floodplain wetlands.

**2a.** Wetlands associated with toe slopes and floodplains of small streams of the Coastal Plain where groundwater discharge is a major contributing input source (mixed hydrological regime: occurs in very narrow part of the groundwater driven complex that is influenced by overbank flooding) with alluvial landform a minor part of the complex; smaller order stream floodplain margins where groundwater input also contributes to overall hydrology. These areas are generally small features along streams and are usually not as well-developed as seepage swamps in larger stream systems.....**COASTAL PLAIN FLOODPLAIN** HGM Class: Riverine or Slope

**2b.** Wetlands associated with distinct depressional and slope geomorphic features.

**3a.** Basin wetlands, depressions, or very flat areas with evidence of ponded water, unidirectional flow not evident, lacks natural outlet, maintained by high water tables and seasonal precipitation. Hydrologic regimes range from saturated to seasonally flooded.

**4.** Seasonally flooded to saturated forested flats and depressions of broad coastal plain terraces (i.e., “wet flatwoods”) with fluctuating water levels and intermittently ponded depressions. Soils are silt, sand, and clay loams, sometimes with a thin (< 30 cm [12 in]) mantle of coarse, fibric peat.

**5a.** Located on flat terraces and shallow depressions with seasonally perched water tables and braided channels...**COASTAL PLAIN FLATWOOD AND DEPRESSION SWAMP** Flatwood: HGM Class- Flat; Depression Swamp: HGM Class- Depression

**5b.** Small (<0.1 ha- 2 ha) shallow pools with a well-defined, discrete basin overlying a clay hardpan or other impermeable soil or rock layer impeding drainage, may or may not have vegetation in basin.....**VERNAL POOL** HGM Class: Depression

**3b.** Slope wetlands associated with groundwater discharge zones (i.e., seeps, springs) and perennial, unidirectional flow towards a natural outlet such as a stream.

**6a.** Small (usually <1m<sup>2</sup>), localized area of groundwater discharge, point source, rare in Coastal Plain.....**SPRING** HGM Class: Slope

**6b.** Larger wetland systems with diffuse drainage patterns, widespread.

**7a.** Open wetlands characterized by predominately shrub and herbaceous vegetation and localized groundwater discharge zones. (*note. Lack of natural disturbances [e.g., fire, beaver activity, grazing] in these habitats often promote woody plant succession.*) Saturated “bog-like” wetlands along gently sloping headwater streams, seepage toe-slopes, and oligotrophic spring-heads with considerable accumulation of peat mosses (*Sphagnum spp.*) at varying depths, soils acidic and infertile (*note. The term “bog” applied here is a technical misnomer since none of these wetland systems in Maryland are ombrotrophic.*).....**COASTAL PLAIN SEEPAGE BOG AND FEN** HGM Class: Organic Soil Flat; Slope

**7b.** Saturated forests of sloping stream headwaters, large spring seeps, lateral seeps in ravines and stream bottoms with diffuse drainage patterns. Braided stream channels, muck-filled depressions, and hummock-and-hollow microtopographic features evident.....**COASTAL PLAIN SEEPAGE SWAMP** HGM Class: Slope or Riverine

Species by vegetation stratum that represent those with high constancy values (>75%) for the more common finer community types (i.e., association level) of Key Wildlife Habitats. Indicator species are those with a high diagnostic value to type, high fidelity, and high relative cover.

Key Wildlife Habitat	Trees	Shrubs	Herbs	Vines	Indicator
<b>Coastal Plain Floodplain</b>	<i>Platanus occidentalis, Liquidambar styraciflua, Liriodendron tulipifera, Quercus michauxii, Fraxinus pennsylvanica, Betula nigra, Acer rubrum</i>  <i>Additional for Blackwater Streams: Taxodium distichum, Nyssa sylvatica, Chamaecyparis thyoides (rare)</i>	<i>Lindera benzoin, Asimina triloba, Ilex opaca, Ilex verticillata, Carpinus caroliniana</i>	<i>Arauptela (Thelypteris) noveboracensis, Mitchella repens, Arisaema triphyllum, Boehmeria cylindrica, Saururus cernuus, Cinna arundinacea, Galium circaezans, Medeola virginiana, Thalictrum thalictroides, Impatiens capensis, Glyceria striata</i>	<i>Toxicodendron radicans, Parthenocissus quinquefolia, Campsis radicans</i>	<i>Platanus occidentalis, Betula nigra, Arauptela (Thelypteris) noveboracensis, Saururus cernuus, Cinna arundinacea</i>  <i>Additional for Blackwater Streams: Taxodium distichum, Nyssa sylvatica, Chamaecyparis thyoides (rare)</i>
<b>Coastal Plain Flatwood and Depression Swamp</b>	<i>Quercus phellos, Quercus palustris, Quercus michauxii, Quercus pagoda, Liquidambar styraciflua</i>	<i>Eubotrys racemosa, Vaccinium corymbosum, Clethra alnifolia</i>	<i>Woodwardia areolata, Osmunda cinnamomea, Mitchella repens, Osmunda regalis, Chasmanthium laxum</i>	<i>Smilax rotundifolia</i>	<i>Quercus pagoda, Quercus michauxii, Vaccinium corymbosum</i>
<b>Coastal Plain Seepage Bog and Fen</b>	<i>Nyssa sylvatica, Acer rubrum, Pinus rigida</i>	<i>Rhododendron viscosum, Toxicodendron vernix, Rubus hispidus, Ilex glabra, Clethra alnifolia</i>	<i>Carex atlantica, Andropogon glomeratus, Rhynchospora gracilentia, Eupatorium pilosum, Dichanthelium dichotomum var. dichotomum</i>	<i>Smilax pseudochina</i>	<i>Smilax pseudochina, Pinus rigida, Andropogon glomeratus, Rhynchospora gracilentia</i>
<b>Coastal Plain Seepage Swamp</b>	<i>Nyssa sylvatica, Acer rubrum, Magnolia virginiana</i>	<i>Clethra alnifolia, Viburnum nudum, Rhododendron viscosum</i>	<i>Woodwardia areolata, Osmunda cinnamomea, Osmunda regalis, Carex folliculata, Symplocarpus foetidus</i>	<i>Smilax rotundifolia</i>	<i>Magnolia virginiana, Clethra alnifolia, Viburnum nudum, Symplocarpus foetidus</i>

**Vernal Pools and Springs** have limited to sparse herbaceous and/or shrub vegetation in the wetland basin. Some Springs have *Sphagnum* species. The surrounding vegetation will represent one of the KWH listed here. Vernal Pools and Springs are most likely to be embedded in Coastal Plain Floodplain, Coastal Plain Flatwood and Depression Swamp, or Coastal Plain Seepage Swamp.

#### **SOIL/SUBSTRATE (Section 4.4)**

Healthy soil function supports plant life and biogeochemical processing for nutrient storage and transformation. Surface features such as changes in elevation over a small area (microtopography) can add to the complexity of the habitat and increase biodiversity, and organic matter accumulation and nutrient dynamics are influenced by leaf litter and ground cover. Disturbance of the surface layer increases the potential for erosion or sedimentation. Prior to fieldwork, mapped soil characteristics for the site should be reviewed. Note any deviations from these characteristics on the data sheet as well as indications of soil compaction and disturbances. Depth to water table and/or extensive roots in the soil should be noted on the data sheet. Examine a soil sample to determine all of the standard measures on the data sheet unless the floodplain does not naturally have hydric soils and/or does not have functioning hydric soils under current conditions. In that case, only score Microtopography, Organic Matter Accumulation, and Soil Disturbance. Note the presence of a gravelly substrate in the Observations/Comments section on the data sheet.

**Redox Concentrations** -Do not score if the floodplain does not naturally have hydric soils and/or does not have functioning hydric soils under current conditions (e.g., relict conditions). Consider depth to groundwater and if other water sources are altered or still sufficient to contribute to reducing conditions. **Extract a sample that is 18" deep from a representative area of the AA** where the soil has not obviously been disturbed. You may need to break open the soil sample to effectively see the rusty red redox concentrations. See Manual for guidance related to scoring soils with red parent material or other problematic soils.

Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Biogeochemical cycling excellent, with redox concentrations starting 0 to 6" from the soil surface and covering >10% of the surface area.	
Good = 3	Biogeochemical cycling good, with redox concentrations starting >6" to 12" from the soil surface and covering >10% of the surface area OR redox concentrations start 0-6" from the soil surface and represent <10% of the surface area.	
Fair = 2	Biogeochemical cycling fair, with redox concentrations starting >12" to 18" from the soil surface and covering >10% of the surface area OR redox concentrations start >6" to 12" from the soil surface and represent <10% of the surface area.	
Poor = 1	Biogeochemical cycling poor, with redox concentrations starting >12" to 18" from the soil surface and covering <10% of the surface area OR no redox concentrations within 18" of the soil surface.	

**Soil Organic Matter**- Do not score if the floodplain does not naturally have hydric soils and/or does not have functioning hydric soils under current conditions. Consider depth to groundwater and if other water sources are altered or still sufficient to contribute to reducing conditions. **Examine the extracted soil sample** for an organic surface horizon or determine features of the mineral surface layer(s).

Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Organic surface horizon present (any thickness).	
Good = 3	Mineral surface layer(s) are $\geq 4$ " thick with matrix value $\leq 3$ and chroma $\leq 2$ .	
Fair = 2	Mineral surface layer(s) are $< 4$ " thick with matrix value $\leq 3$ and chroma $\leq 2$ .	
Poor = 1	Mineral surface layer(s) are $< 4$ " thick with matrix value $> 3$ and $\leq 4$ or chroma $> 2$ and $\leq 3$ .	

**Microtopography**- Estimate the percent of the AA with an elevation change of at least 3" due to soil elevations and woody debris in an advanced stage of decomposition. Microtopography is often present as vegetated hummocks, raised areas that support tree trunks and roots, or nursery logs.

Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	More than 50% of the AA shows at least a 3" increase in elevation over the base elevation of the AA.	
Good = 3	30-49% of the AA shows at least a 3" increase in elevation over the base elevation of the AA.	
Fair = 2	10-29% of the AA shows at least a 3" increase in elevation over the base elevation of the AA.	
Poor = 1	<10% of the AA shows at least a 3" increase in elevation over the base elevation of the AA.	

**Organic Matter Accumulation- Organic Matter Accumulation**- Indicators will vary with season and KWH. Estimate the percent cover of herbaceous and woody plants, both living and dead residue. **Estimate how much of the AA is covered by >1" of loose leaf litter OR by at least 5 stacked layers of decaying or wetted leaves.** When leaf litter depth is naturally lower, pick apart decaying or wetted leaves to determine if there are 5 or more stacked layers and estimate percent coverage.

Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Organic matter accumulation from root turnover/leaf litter is high as herbaceous and woody plant ground cover plus leaf litter covers >75% of the surface. To count towards coverage, loose leaves must be at least 1" thick or decaying leaves must have at least 5 stacked layers.	
Good = 3	Organic matter accumulation from root turnover/leaf litter is moderate as herbaceous and woody ground cover plus leaf litter covers >50-74% of the surface. To count towards coverage, loose leaves must be at least 1" thick or decaying leaves must have at least 5 stacked layers.	
Fair = 2	Organic matter accumulation from root turnover/leaf litter is low as herbaceous and woody ground cover plus leaf litter covers >25-50%. To count towards coverage, loose leaves must be at least 1" thick or decaying leaves must have at least 5 stacked layers.	
Poor = 1	Organic matter accumulation from root turnover/leaf litter is minimal as herbaceous or woody ground cover plus leaf litter covers $\leq 25\%$ . To count towards coverage, loose leaves must be at least 1" thick or decaying leaves must have at least 5 stacked layers.	

**Soil Disturbance**- Note impacts to the soil surface as indicated by bare soil, unless caused by natural factors or the soil is naturally bare. Look at the extent of impact across the AA and the greatest depth of the impact (including ponding or channeling of water).

Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Little bare soil OR bare soil and soil disturbed areas are limited to naturally caused disturbances such as flood deposition, game trails, beaver activity, etc. OR soil is naturally bare. No human-caused impacts evident.	
Good = 3	Minor amounts or localized, small patches of bare or disturbed soil are present from factors such as cattle trampling or heavy grazing that leads to erosion, compaction or trampling by machinery, ruts or other disturbances from ATV or other vehicular activity, sedimentation due to	

	human causes, or invasive earthworms. Extent of impact is minimal and greatest depth is limited to a few centimeters (a few inches) and does not show evidence of ponding or channeling of water.
Fair = 2	Moderate amounts of bare or disturbed soil are present due to human-caused activities. Extent of impact is moderate and greatest depth may extend 5–10 cm (2–4 inches), with localized deeper ruts. Shows some evidence of ponding or channeling of water.
Poor = 1	Substantial amounts of bare or disturbed soil are present due to human-caused activities. Impact is extensive with long-lasting impacts. Greatest depth of impact extends > 10 cm (4 inches); deeper ruts may be widespread and show some evidence of extensively altering hydrology (e.g., ponding or channeling of water).

### HYDROLOGY (Section 4.5)

Hydrology is a complicated ecological factor to measure during a rapid assessment, as the evaluation of one metric partly relates to another. In this section, two aspects of the hydrology of the AA are scored by indicating the presence of natural and altered features of the Water Source and Hydroperiod and Hydrologic Connectivity. The scoring for these metrics varies depending on the type of KWH, so make sure you are using the correct scoring table. The Stream Bank and Channel metric, in contrast, is assessed for the entire project area using indicators of alteration as well as stabilization and recovery. Data sheet check boxes will capture features for scoring mentioned in the following sections. Obstructions, alterations, and point source discharges may be visible on aerial photos or other available imagery. LiDAR Hillshade images may assist with identifying existing channels and other relevant features.

**Water Source (Section 4.5.1)** This metric focuses on the forms and places of direct inputs of water to the AA, as well as any unnatural diversions of water from the AA or other features that affect saturation of the wetland. **Focus on the main source of water for this evaluation and use the scoring table for the correct KWH.** Note evidence of natural and unnatural/manipulated characteristics using the check boxes on the data sheet. Consider whether alterations are recent and if they are currently having a negative effect. Beaver activity, although it may have caused changes, should be considered as a natural change for scoring.

<b>Coastal Plain Floodplain:</b> Groundwater discharge not a major input. For scoring, note stream bank alterations that will affect the riparian water source.		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Water source is natural. Lacks point charge discharges into or adjacent to the site. No unnatural obstructions to water source or impact on overland flow and overbank flooding. Plant community reflective of characteristic KWH or not altered by natural changes to water source.	
Good = 3	Water source is mostly natural, but wetland directly receives occasional or small amounts of inflow from anthropogenic sources such as some road runoff, small storm drains, or other minor point source discharges emptying into the wetland. Up to 25% of stream banks are affected due to dikes, rip rap and/or elevated culverts, or there is increased discharge due to other causes. Little change in plant community resulting from unnatural alterations.	
Fair = 2	Water sources are moderately impacted by anthropogenic sources but are still a mix of natural and non-natural sources. Between 25-75% of stream banks are affected (e.g., dikes, rip rap, concrete, and elevated culverts) or increased discharge due to other causes. Wetlands still present due to groundwater or other water inputs, but potentially reduced in extent and showing some plant community changes; or plant community changes due to increased unnatural water inputs.	
Poor = 1	Water source contains a substantial amount of inflow from anthropogenic sources, such as major point source discharges into or adjacent to the wetland. > 75% of stream banks are affected (for example due to dikes, rip rap, concrete, and elevated culverts) or increased discharge due to other causes. Wetlands are reduced in extent unless high groundwater or other surface water inputs maintain them. Plant community changes are observed due to unnatural water inputs.	
<b>Coastal Plain Floodplain:</b> Mixed hydrologic regime with some input from groundwater and from precipitation or limited flooding		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Water source is natural. Lacks point charge discharges into or adjacent to the site. No unnatural obstructions to lateral or vertical movement of ground or surface water. Plant community reflective of characteristic KWH or not altered by natural changes to water source.	
Good = 3	Water source is mostly natural, but wetland directly receives occasional or small amounts of inflow from anthropogenic sources such as some road runoff, small storm drains, or other minor point source discharges emptying into the wetland. Minor restrictions to the lateral or vertical movement of ground or surface waters by unnatural features. Little change in plant community resulting from unnatural alterations.	
Fair = 2	Water sources are moderately impacted by anthropogenic sources, but are still a mix of natural and non-natural sources. Wetland is still connected to its natural water source (e.g., modified ponds on a floodplain that are still connected to alluvial aquifers, natural stream channels that now receive substantial irrigation return flows, many small/few large storm drains), but moderately disconnected from floodplain due to multiple geomorphic modifications. Moderate restrictions to the lateral or vertical movement of ground or surface waters by unnatural features. Wetlands still present due to groundwater or other water inputs, but limited reduction in extent and showing some plant community changes; or some limited plant community changes due to increased unnatural water inputs.	
Poor = 1	Water source contains a substantial amount of inflow from anthropogenic sources, such as major point source discharges into or adjacent to the wetland. Wetland has reduced connection to natural water source (e.g., loss of overbank flow). Wetlands are potentially reduced in extent if no other surface water inputs maintain them. Plant community changes are observed due to unnatural water inputs.	

<b>All other KWH:</b> Predominantly groundwater or precipitation water source, with potential limited flooding from small stream in relation to wetlands in riparian system		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Water source is natural. Lacks point charge discharges into or adjacent to the site. Groundwater or precipitation dominant or only water source; otherwise, no unnatural obstructions to lateral or vertical movement of ground or surface water, or, if perched water table, impermeable soil layer is intact. Plant community reflective of characteristic KWH or not altered by natural changes to water source.	
Good = 3	Water source is mostly natural, but wetland directly receives occasional or small amounts of inflow from anthropogenic sources such as some road runoff, small storm drains, or other minor point source discharges emptying into the wetland. Minor restrictions to the lateral or vertical movement of ground or surface waters by unnatural features, such as levees or excessively high banks (less than 25% of the site). If perched, impermeable soil layer partly disturbed. Little change in plant community resulting from water source alterations.	
Fair = 2	Water source is moderately impacted by anthropogenic sources, but still a mix of natural and non-natural sources. Moderate restrictions to the lateral or vertical movement of ground or surface waters by unnatural features or alteration. Between 25-75% of the site is restricted by barriers to drainage. If perched, impermeable soil layer moderately disturbed. Drainage back to the wetland is incomplete due to impoundment. Wetlands still present due to groundwater or other water inputs, but limited reduction in extent and showing some plant community changes; or some limited plant community changes due to water source alterations.	
Poor = 1	Water source contains a substantial amount of inflow from anthropogenic sources, such as major point source discharges into or adjacent to the wetland. Most or all water stages are contained within artificial banks, levees, or comparable features. Greater than 75% of wetland is restricted by barriers to drainage. If perched, impermeable soil layer strongly disturbed. Wetlands reduced in extent and show plant community changes due to water source alterations.	

**Stream Bank and Channel (Section 4.5.2)** Indicate the characteristics of the stream bank and channel for the project area using the check boxes on the data sheet and additional lines as needed, including evidence of equilibrium, signs of recovery, channel and bank instability and their sources. **This score will apply to all AA in the project area.** Examples of field indicators of equilibrium, degradation, and aggradation are presented in the table on the next page. If available, indicate the Bank Erosion Hazard Index (BEHI) score, Near Bank Stress (NBS) score, and modeled inundation from storm events and use them in your scoring process. Use online resources (Section 3.1) to fill in the Benthic Index of Biotic Integrity (IBI) and Fish IBI Values and Ratings if available.

<b>Stream Bank and Channel in Project Area (score applies to all AA in project area)</b>		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Indicators of channel equilibrium present. Minimal or no evidence of degradation or aggradation leading to channel instability or migration. Bank instability none or minimal. Channel is not unnaturally entrenched. If calculated, BEHI/NBS scores low.	
Good = 3	Minor channel incision. Channel is somewhat entrenched (overbank flow occurs during most floods). Some evidence of degradation or aggradation leading to a minimal level of channel instability or migration. Minor bank instability. If calculated, BEHI/NBS scores low.	
Fair = 2	Channel is incised. Channel is moderately entrenched (overbank flow only occurs during moderate to severe floods, functioning at risk). Uncharacteristic aggradation or degradation is present leading to a moderate level of channel instability or migration. Bank instability moderate. BEHI/NBS scores moderate.	
Poor = 1	Channel is incised. Channel is substantially entrenched (overbank flow never occurs or only during severe floods-not functioning). Channel entirely or extensively disconnected from the floodplain. Bank instability substantial. BEHI/NBS scores high, very high, or extreme.	

**Hydroperiod and Hydrologic Connectivity (Section 4.5.3)** This metric examines the characteristic frequency, level, and duration of wetland inundation or saturation, regardless of the source, and the ability of water to flow into or out of the wetland. **Use the scoring table for the correct KWH and check off what you observe on the data sheet.** Estimate the hydroperiod variation based on visual indicators and soil redox. Indicators of changes in extent and duration of inundation or saturation are presented on the next page. If available, add information for storm interval flooding, Bank Height Ratio, and Entrenchment Ratio.

<b>Coastal Plain Floodplain Note:</b> Recent beaver activity may lead to deviations from rating descriptions. This should be noted on the data sheet.		
Low natural variation of hydroperiod _____ High natural variation of hydroperiod _____		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Evidence of recent overbank flooding. Completely connected to floodplain (backwater sloughs and channels, may be still pools with flow). No major hydrologic stressors present that impact natural hydroperiod or impact due to natural events (e.g., beaver dams). No unnatural obstructions to lateral or vertical movement of ground or surface water.	
Good = 3	Evidence of overbank flooding. Minimally disconnected from floodplain. Minor alterations in frequency, levels, or duration of hydroperiod. Minor restrictions to the lateral or vertical movement of ground or surface waters by unnatural features. Flooding at 2-year storm interval.	
Fair = 2	Some evidence of overbank flooding, likely during larger storm events. Moderately disconnected from floodplain due to multiple geomorphic modifications. Moderate restrictions to the lateral or vertical movement of ground or surface waters by unnatural features. Moderate flooding at 10-year storm interval.	
Poor = 1	Overbank flooding generally no longer occurs. Disconnected from floodplain, likely causing some drainage of groundwater. Flooding may or may not occur at 100-year or greater storm interval.	

Other KWH		
Low natural variation of hydroperiod _____ High natural variation of hydroperiod _____		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Overbank flooding present and recent but not predominant water source to wetland. No unnatural obstructions to lateral or vertical movement of ground or surface water.	
Good = 3	Evidence of overbank flooding but not predominant water source to wetland. Hydroperiod with minor alterations in frequency, levels, or duration due to groundwater and other inputs. Minor restrictions to the lateral or vertical movement of ground or surface waters by unnatural features.	
Fair = 2	Some evidence of overbank flooding, likely during larger storm events. Hydroperiod with moderate alterations in frequency, levels, or duration due to groundwater and other inputs. Moderate restrictions to the lateral or vertical movement of ground or surface waters by unnatural features.	
Poor = 1	Overbank flooding generally no longer occurs. Hydroperiod with substantial alterations in frequency, levels, or duration due to groundwater and other inputs. Substantial restrictions to the lateral or vertical movement of ground or surface waters by unnatural features.	

Condition	Field Indicators for Stream Bank and Channel and Hydroperiod for Coastal Plain Floodplain
Indicators of Channel Equilibrium	<ul style="list-style-type: none"> <li>• The channel (or multiple channels in braided systems) has a well-defined usual high water line, or bankfull stage, that is clearly indicated by an obvious floodplain. A topographic bench represents an abrupt change in the cross-sectional profile of the channel throughout most of the site.</li> <li>• The usual high water line (consistent with ACOE ordinary high water mark) or bankfull stage corresponds to the lower limit of riparian vascular vegetation.</li> <li>• The channel contains embedded woody debris of the size and amount consistent with what is available in the riparian area.</li> <li>• There is little or no active undercutting or burial of riparian vegetation.</li> </ul>
Indicators of Active Degradation (Erosion)	<ul style="list-style-type: none"> <li>• Portions of the channel are characterized by deeply undercut banks with exposed living roots of trees or shrubs. There are abundant bank slides or slumps, or the banks are uniformly scoured and unvegetated.</li> <li>• Riparian vegetation may be declining in stature or vigor, and/or riparian trees and shrubs may be falling into the channel.</li> <li>• The channel bed lacks any fine-grained sediment (unless it is the dominant bank material).</li> <li>• Recently active flow pathways appear to have coalesced into one channel (i.e., a previously braided system is no longer braided).</li> </ul>
Indicators of Excessive Aggradation (Sedimentation)	<ul style="list-style-type: none"> <li>• The channel through the site lacks a well-defined usual high water line.</li> <li>• There is an active floodplain with fresh splays of excessive sediment covering older soils or recent vegetation.</li> <li>• There are partially buried tree trunks or shrubs.</li> <li>• Excessive cobbles and/or coarse gravels have recently been deposited on the floodplain.</li> <li>• There are partially buried, or sediment-choked, culverts.</li> </ul>
Condition	Hydroperiod Field Indicators for Other KWH Types
Reduced Extent and Duration of Inundation or Saturation	<ul style="list-style-type: none"> <li>• Upstream diversions, impoundments, pumps, ditching, or draining from the wetland.</li> <li>• Water withdrawal (wells).</li> <li>• Evidence of aquatic wildlife mortality.</li> <li>• Encroachment of terrestrial vegetation.</li> <li>• Encroachment of young, tall, vigorous trees if not usually present, shading of underlying mosses.</li> <li>• Stress or mortality of hydrophytes or sphagnum.</li> <li>• Compressed or reduced plant zonation.</li> <li>• Organic soils occur well above contemporary water tables.</li> <li>• Increased discharges resulting in channel downcutting.</li> </ul>
Increased Extent and Duration of Saturation	<ul style="list-style-type: none"> <li>• Berms, dikes, or other water control features that increase duration of ponding (e.g., pumps).</li> <li>• Diversions, ditching, or draining into the wetland.</li> <li>• Late-season vitality of annual vegetation.</li> <li>• Recently drowned riparian or terrestrial vegetation (e.g., beaver-created impoundment).</li> <li>• Extensive fine-grained deposits on the wetland margins.</li> </ul>

## KEY WILDLIFE HABITAT AND VEGETATION COMPOSITION (Section 4.6)

Vegetation structure and composition are of particular interest for assessing the condition of Key Wildlife Habitats because they directly support the ecological needs of animal and plant species of concern. In this section, metrics provide information on the interspersed vegetation patches, habitat features/evidence of animal use, vertical structure, and standing and downed woody debris (standing tree snags and downed trees and branches). Vegetation data collected previously or simultaneously using standard wetland delineation methods are used to document vegetation composition and can be used to assess most metrics. Scores are assigned to reflect the presence and extent of invasive and native plant species in herbaceous and woody layers, including the presence of native species that are diagnostic (Section 4.3) and indicative of disturbance. Additionally, any plant species listed as rare, threatened, or endangered in Maryland should be identified (see Manual for source of current list). **These species should be noted on the data sheet even if they are not dominant.** A Floristic Quality Assessment will be calculated using the Excel data sheet or as otherwise described in the Manual. **Expected conditions vary by Key Wildlife Habitat for some metrics- use the correct scoring tables.**

**Interspersion and Patch Richness (Section 4.6.1)** For this metric, interspersion and patch richness will be scored separately and then averaged for a final score. Interspersion is assessed within the AA but patch richness is assessed within the AA and out to 10m around the AA on each side.

**Interspersion:** The figures below show a range of patterns for the interspersion of vegetation patches for different Key Wildlife Habitats. Different vegetation types, such as hummocks, sphagnum, shrub areas, patches of herbaceous vegetation, and patches or lines of trees of different heights or ages, should be noted for the AA. **Select the diagram below for the appropriate KWH to determine a score for this metric.** To be considered, vegetative patches should represent at least 5% of the AA in single or multiple locations. This metric is often reflective of the topographic complexity metric in many wetland types. Record the score on the next page.

**Coastal Plain Seepage Swamp, Coastal Plain Seepage Bog and Fen, Coastal Plain Flatwood and Depression Swamp, Vernal Pool, Spring.**  
(Source: USACE 2015 Texas Rapid Assessment Method)

**Scoring: High = 4** Vegetation patches are large and intertwined or numerous and scattered

**Moderate = 3** At least two types of vegetation patches are present but patches are slightly smaller or less scattered/intertwined than "High" category

**Low = 2** Two types of vegetation patches are present but in smaller, very localized, and/or isolated patches

**None = 1** Only one type of vegetation patch is present




**Coastal Plain Floodplain:** The red box represents the boundary of the AA and each color represents a unique plant zone such as shrub areas, patches of herbaceous vegetation, or tree clumps of different ages or heights. The speckled background represents the background matrix of vegetation and the blue line represents the stream. For multithread stream systems, evaluate the channel with the highest complexity of plant zones for scoring. (Source: California Rapid Assessment Methods for Wetlands Riverine Wetlands Field Book 2013)

**Scoring: A = 4** High complexity of scattered and intertwined plant zones

**B = 3** Moderate complexity of intertwined plant zones

**C = 2** Minimal complexity of plant zones with little interspersion

**D = 1** Few plant zones with localized, isolated patches



A

B

C

D

**Patch Richness:** Patch richness provides a measure of components that represent potential wildlife habitat. **Check the following features off on the data sheet if they are present in the AA or within 10 m (33 feet) of the AA boundary.** Count the number of features present. Also indicate the presence of any observed wetland- or stream-associated animals such as frogs, waterbirds, crayfish, fish, mussels, etc. on the data sheet. Record the score on the next page.

**Features:** Spring or upwelling groundwater; Depression; Vegetated pool; Unvegetated pool; Unvegetated flat; Island; Animal mound or burrow; Beaver dam or lodge; Beaver-chewed vegetation; Oxbow, swale, secondary channel; Wind-thrown tree hole; Mound;



Bank overhang with tree roots; Tip-up tree root mound; Brush piles; Abundant deciduous leaf litter; Partially buried natural debris; Debris jam; Plant hummock/tussocks; Other wildlife habitat

Score	Coastal Plain Floodplain, Coastal Plain Seepage Bog and Fen, Coastal Plain Seepage Swamp	Coastal Plain Flatwood and Depression Swamp	Vernal Pool/Spring
4	≥ 6	≥ 7	≥ 4
3	5 - 6	6 - 7	3 - 4
2	3 - 4	4 - 5	2
1	< 3	< 4	< 2

**Interspersion and Patch Richness Score:** Calculate the mean of the Interspersion and Patch Richness metrics below. Use the table to assign an overall score for this metric.

Score	Mean of Interspersion and Patch Richness Metric Scores
Excellent = 4	3.5 – 4
Good = 3	2.6 - 3.4
Fair = 2	1.6 – 2.5
Poor = 1	1 – 1.5

**Interspersion Score:** \_\_\_\_\_

**Patch Richness Score:** \_\_\_\_\_

**Mean of Interspersion and Patch Richness Scores:** \_\_\_\_\_

**Overall Score for Metric (see table at left):** \_\_\_\_\_

**Vertical Structure (Section 4.6.2)** This metric provides an assessment of the overall structural complexity of vegetation layers, including presence of multiple strata, age and structural complexity of canopy layer, and effects of disease or mortality on structure. **Assess within the AA and out to 10m (33 feet) of the AA boundary.** Forested KWH are assessed differently than non-forested KWH (Coastal Plain Seepage Bog and Fen). As **beaver activity** can impact vertical structure, the vertical structure in the surrounding area and previous structure as indicated by snags and downed trees should be considered when assigning a score. Note the presence of these changes on the data sheet. **Vernal Pools and Springs** are expected to have only sparse woody and/or herbaceous vegetation in the basin area, if any. For these KWH, assess the vertical structure in the surrounding area. For **Coastal Plain Seepage Bog and Fen**, an evaluation of the integrity of dominant growth forms is made (e.g., whether shrubs have been removed, killed, or increased or if the herbaceous layer has been reduced or homogenized by stressors). Reference to the description for this KWH can be useful. **Use the correct KWH table and assign the rating to the category with the majority of features present.**

Coastal Plain Floodplain, Coastal Plain Flatwood and Depression Swamp, Coastal Plain Seepage Swamp		
<b>Vernal Pool and Spring:</b> only assess structure in area surrounding basin- limited to sparse herbaceous vegetation is usually present in the basin area. <b>Note:</b> Recent beaver activity may lead to deviations from rating descriptions for Coastal Plain Floodplain. This should be noted on the data sheet.		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Tree canopy or highest woody level present is a heterogeneous mosaic of patches of different ages or sizes. Gaps of varying size. Multiple layers are created through the presence of trees of varying ages and heights and the shrub layer. Large trees (> 60 cm or 24" dbh) expected to be present (≥ 10% of trees present). If large trees are absent, few or no large stumps are present and there is evidence of a natural disturbance event (e.g., large downed wood from wind storms, fire scars, beaver activity, tree senescence). Little impact from deer browse.	
Good = 3	Tree canopy or highest woody level present is largely heterogeneous in age or size. Multiple layers are present, but one layer missing or little variation in ages and heights of woody vegetation in at least one layer. Less than 10% of trees present are large trees (>60 cm or 24" dbh) due to human activities. At least 20% of trees present are >30 cm or 12" dbh. Minor presence of cutting, browsing, grazing and other degradation such as forest pest/pathogens. If large trees are absent, few or no large stumps are present and there is evidence of a natural disturbance event (e.g., large downed wood from wind storms, fire scars, beaver activity, tree senescence). Little impact from deer browse.	
Fair = 2	Tree canopy or highest woody level present is somewhat homogeneous in age or size. More than one layer present, but one or more layers missing. Little variation in ages and heights of woody vegetation in layers. Less than 20% of trees present are >30 cm or 12" dbh are present. Moderate levels of cutting, browsing, or grazing, or other degradation such as forest pest/pathogens has caused the loss of larger trees rather than a natural disturbance event.	
Poor = 1	Tree canopy or highest woody level present is very homogeneous in age or size. Only one or two layers present due to human activities. Most, if not all, larger trees (dbh 30-60 cm or 12-24") have been removed. Major cutting, heavy browsing, grazing, or other degradation such as forest pest/pathogens.	
Coastal Plain Seepage Bog and Fen		
Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Mortality of woody vegetation, if present, is due to natural factors such as wind storms or senescence. Excellent potential for site recovery given structure present and lack of degradation (past or present). <b>Bogs/acidic fens:</b> Peatland structure includes shrub and herb strata (some tall and some short). When present (peatland not too wet), trees are relatively short and stunted with rounded tops and furrowed bark. Shrubs are < 50 cm and open enough to allow for a nearly continuous ground cover of <i>Sphagnum</i> and other expected vegetation around tree/shrub bases AND in low hummocks, hollows, or other low areas.	

	<u>Circumneutral/rich fens</u> : Primarily short-statured vegetation and nearly continuous cover of mosses (except in tall sedge fens - which are naturally more vigorous, homogenous, and often with little bryophyte cover). Shrubs may be present as a mosaic with open areas. Tree species, when present, do not form a closed canopy. <i>Sphagnum</i> and other mosses actively growing. Never more than local, small patches of degenerating <i>Sphagnum</i> .
Good = 3	Minor negative anthropogenic influences present, or the site is still recovering from major past human disturbances. Mortality or degradation due to grazing, peat mining, limited timber harvesting, or other anthropogenic factors may be present, though not widespread. The site can be expected to meet minimally disturbed conditions in the near future if negative influences do not continue. <u>Bogs/acidic fens</u> : Shrubs and herbs show minor alterations from expected conditions. A few areas of dense and tall shrubs (> 1 m) may occur (dense enough to eliminate <i>Sphagnum</i> /moss growth). Some trees may have been or killed due to anthropogenic stressors. <u>Circumneutral/rich fens</u> : Shrubs and herbs show minor alterations from expected conditions.
Fair = 2	Expected structural classes are not present. Shrubs and herbs moderately altered from expected conditions. The site will recover to minimally disturbed conditions only with the removal of degrading influences and moderate recovery times. <u>Bogs/acidic fens</u> : Shrub cover averages > 1 m tall and is beginning to reduce <i>Sphagnum</i> cover. Many trees have been cut or killed due to anthropogenic stressors. <u>Circumneutral/rich fens</u> : Trampling or other physical disturbance has moderately reduced moss cover where expected. Overall, evidence of degradation includes moderate levels of cutting, mowing, browsing, fire or grazing. <i>Sphagnum</i> still regenerating in open areas.
Poor = 1	Expected peatland structure is absent or much degraded due to anthropogenic factors, such as peat mining. Overall, evidence of degradation includes major cutting, mowing, browsing, fire or grazing. Woody regeneration is minimal and existing structure is in poor condition, unnaturally sparse, or depauperate. Shrubs and herbs substantially altered from expected conditions. Recovery to minimally disturbed condition is questionable without restoration, or will take many decades. <u>Bogs/acidic fens</u> : Most if not all <i>Sphagnum</i> cover has been eliminated due to extremely dense and tall (> 1 m) shrubs. Trees have all been cut or killed by anthropogenic stressors. <u>Circumneutral/rich fens</u> : Trampling or other physical disturbance has eliminated moss cover where it is expected. <i>Sphagnum</i> not regenerating, even in open areas.


**Standing and Downed Woody Debris (Section 4.6.3)** Standing or fallen woody debris (snags and downed branches and trees) plays a critical role in riparian systems. Estimation of coarse woody debris should be based on a walkthrough of the entire AA if possible. For large AA, estimation along transects may be preferred. Use the check boxes in the data sheet to indicate features present for the correct KWH. In forested KWH, pay special attention to the amount of coarse woody debris when surveying the AA and note the creation of woody debris from cutting, pests/pathogens, or other factors. Riverine wetlands that have incised banks, no longer experience flooding, experience overgrazing, or are no longer at a dynamic equilibrium may lack coarse woody debris. For wetlands dominated by shrub and herb layers, note the quantity and distribution of litter compared with the baseline that may be expected in the landscape. Active floodplain systems are typically low in litter. As **Vernal Pools and Springs** may have only scattered woody debris, evaluate both the basin and the surrounding area. Peatlands are dominated by peat-forming species which contribute enough litter and debris to maintain carbon dynamics, playing a critical role in these systems that may naturally include little coarse woody debris.

<b>Coastal Plain Floodplain, Coastal Plain Flatwood and Depression Swamp, Coastal Plain Seepage Swamp</b>		
<b>Vernal Pool and Spring:</b> assess presence in immediate surrounding area as well as the basin. If non-natural sources have created standing and/or downed woody debris, indicate this on the data sheet.		
<b>Score</b>	<b>Assign rating to category with majority of features present:</b>	<b>SCORE _____</b>
Excellent = 4	Wide diversity of sizes for both standing and downed logs, including larger sizes [ $> 30$ cm (12 in) diameter and $> 2$ m (6 ft) long]] present with 5 or more snags per ha (2.5 ac), but not excessive numbers (suggesting disease or other problems). Downed logs are in various stages of decay, from sound and intact to soft pieces that no longer maintain their shape.	
Good = 3	Moderate diversity of sizes for both standing and downed logs, but larger sizes [ $> 30$ cm (12 in) diameter and $> 2$ m (6 ft) long]] are rare. Larger size class present with 2-4 snags per ha, or an increased but not excessive number of snags (suggesting disease or other problems). Downed logs are in various stages of decay, with few soft pieces that no longer maintain their shape.	
Fair = 2	Moderate-low diversity of sizes for both standing and downed logs, but larger sizes [ $> 30$ cm (12 in) diameter and $> 2$ m (6 ft) long]] very rare or not present. Larger size class present with 1-2 snags per ha, or moderately excessive numbers (suggesting disease or other problems). Downed logs are in various stages of decay, but few to no soft pieces that no longer maintain their shape.	
Poor = 1	Low diversity of sizes for both standing and downed logs. Larger size class [ $> 30$ cm (12 in) diameter and $> 2$ m (6 ft) long]] present with $< 1$ snag per ha, or very excessive numbers (suggesting disease or other problems). Downed logs are mostly in early stages of decay.	
<b>Coastal Plain Seepage Bog and Fen</b>		
<b>Score</b>	<b>Assign rating to category with majority of features present:</b>	<b>SCORE _____</b>
Excellent = 4	Typical of the system. Woody vegetation mortality is due to natural factors. Peat accumulation appears to be stable or actively growing. <u>Bogs/acidic fens</u> : <i>Sphagnum</i> is nearly continuous and growing around tree/shrub bases AND in low hummocks, hollows, or other low areas. <u>Circumneutral/rich fens</u> : Dominant species are active peat-formers.	
Good = 3	Minor alterations to system present. <u>Bogs/acidic fens</u> : Mortality or degradation of peat surface due to grazing, limited timber harvesting, anthropogenic fire or other anthropogenic factors may be present, but not widespread. <u>Circumneutral/rich fens</u> : Mortality or degradation of peat surface due to grazing, limited timber harvesting, anthropogenic fire or other anthropogenic factors may be present, but not widespread.	

Fair = 2	Moderate alterations to system present. <u>Bogs/acidic fens:</u> Ground cover has as much bare peat as <i>Sphagnum</i> cover, or nearly so. <u>Circumneutral/rich fens:</u> Dominance of active peat-formers is being reduced in favor of non-peat-forming grasses and forbs.
Poor = 1	Substantial alterations to system present. <u>Bogs/acidic fens:</u> Ground cover is almost all bare peat with very little <i>Sphagnum</i> cover. <u>Circumneutral/rich fens:</u> Cover of active peat-formers dramatically reduced and site is now dominated by non-peat-forming grasses and forbs.

**Vegetation Composition (Section 4.6.4)** Vegetation of the AA is characterized using the five strata version of the wetland delineation determination (USACE 2010). The species composition is assessed relative to the species expected in each stratum for the KWH. **The coverage of invasive species and native species (both diagnostic and those indicative of disturbance) should be noted regardless of percent cover.** These species are listed with Section 4.3 above. State rare species should be noted. In addition, the sources of stressors or alterations to the native plant community should be noted on the data sheet as well as suggestions for improving native species cover. The diagrams below may be useful to assist with the estimation of percent cover.

**% Cover Estimation Diagrams** (*johnmuirlaws.com and Terry and Chilingar 1955*)



**Invasive Species (Section 4.6.5)** Invasive species are non-native species that can spread into natural ecosystems, where they can displace native species and cause major alterations to KWH. The most common plant invasive species in Coastal Plain stream-associated wetlands are *Microstegium vimineum*, *Glechoma hederacea*, *Rosa multiflora*, *Ligustrum sinense*, *Lonicera japonica*, *Phalaris arundinacea*, and *Phragmites australis*. Identification references and additional species can be found in the Manual. Scoring for Vernal Pools and Springs should use observations from the basin and surrounding area, as only limited sparse vegetation may be present in the basin.

<b>Coastal Plain Floodplain, Coastal Plain Flatwood and Depression Swamp, Coastal Plain Seepage Swamp, Coastal Plain Seepage Bog and Fen</b>		
<b>Vernal Pool and Spring:</b> assess vegetation structure in area surrounding basin, as only limited to sparse vegetation may be present in the basin area.		
<b>Score</b>	<b>Assign rating to category with majority of features present:</b>	<b>SCORE _____</b>
Excellent = 4	Invasive species are absent from all layers or absolute cover in any one woody layer (if present) and herbaceous layer is <1%.	
Good = 3	Invasive species are sporadic (no more than 1-5% absolute cover in any layer).	
Fair = 2	Absolute cover of Invasive species is >5-10% in any one woody layer (if present) and/or present with moderate absolute cover (>5-30%) in the herbaceous layer. Patches of native vegetation are reduced in size and complexity due to the presence of invasive species.	
Poor = 1	Absolute cover of Invasive species is over 10% in any one woody layer (if present) and/or is very abundant (over 30%) in the herbaceous layer. Vegetation reduced in size and complexity due to human disturbance. Patches of native vegetation are reduced in size and complexity due to the presence of invasive species.	

**Native Species (Section 4.6.6)** The presence and composition of native plant species provides an indication of KWH ecological integrity and how well the AA supports a diversity of native animal species. This metric uses the presence of indicator species and characteristic native species for the KWH in the AA (Section 4.3) as well as the presence of native species that indicate human disturbance. Metrics are adjusted for Coastal Plain Seepage Bog and Fen systems and some Spring KWH due to the importance of *Sphagnum*. Indicate the stressors present in the AA on the data sheet and provide suggestions for improvement.

**Native Species Indicative of Disturbance:** These species are those that seem to be more or less weedy and not picky about habitat, or they occur in young, often heavily altered wetland communities. Note the presence of these species to help assess the site and to assist with scoring.

<i>Phalaris arundinacea</i>	<i>Dichanthelium boscii</i>
<i>Typha latifolia</i>	<i>Dichanthelium sphaerocarpon</i>
<i>Elymus glabriflorus</i>	<i>Paspalum floridanum</i>
<i>Muhlenbergia schreberi</i>	<i>Echinochloa muricata</i>
<i>Carex blanda</i> , <i>C. frankii</i>	<i>Coleataenia anceps</i>
<i>Dichanthelium scoparium</i>	<i>Panicum dichtomiflorum</i>

**Coastal Plain Floodplain, Coastal Plain Flatwood and Depression Swamp, Coastal Plain Seepage Swamp, Coastal Plain Seepage Bog and Fen** (see information on second and third pages for diagnostic native species and those that indicate disturbance)

**Vernal Pool and Spring:** assess vegetation structure in area surrounding basin, as only limited to sparse vegetation is usually present in the basin area.

**Note:** Recent beaver activity may lead to deviations from rating descriptions for Coastal Plain Floodplain. This should be noted on the data sheet and considered in assignment of the score.

Score	Assign rating to category with majority of features present:	SCORE _____
Excellent = 4	Herbaceous and woody layers (if present) dominated by diagnostic native species. Layers may be sparse and patchy in areas with deeper flooding, with patches of vegetation confined to hummocks. In other areas, diverse native vegetation present unless there has been a recent natural disturbance.  <u>Bog and Fen, some Springs:</u> <i>Sphagnum</i> is nearly continuous and growing around tree/shrub bases AND in low hummocks, hollows, or other low areas.	
Good = 3	Some diagnostic native species absent or substantially reduced in abundance OR low cover (<10%) of native species indicative of human disturbance. Layer may be sparse and patchy in areas with deeper flooding.  <u>Bog and Fen, some Springs:</u> <i>Sphagnum</i> and other mosses actively growing, but may be eliminated from some areas due to disturbance or invasive species.	
Fair = 2	Few diagnostic species are present. Native species indicative of human disturbance are present with moderate cover (10-30%). Patches of native vegetation are reduced in size and complexity due to human disturbance.  <u>Bog and Fen, some Springs:</u> <i>Sphagnum</i> cover reduced but still regenerating in open areas. Dominance of active peat-formers is being reduced in favor of non-peat-forming grasses and forbs.	
Poor = 1	Few to no diagnostic species are present. Native species indicative of human disturbance are present with >30% cover. Patches of native vegetation are reduced in size and complexity due to human disturbance.  <u>Bog and Fen, some Springs:</u> Very little <i>Sphagnum</i> cover. Cover of active peat-formers dramatically reduced and site is now dominated by non-peat-forming grasses and forbs.	

#### Floristic Quality Assessment (Section 4.6.7)

This method derives an estimate of nativity or habitat quality based on a combination of the tolerance to disturbance or environmental stress and the fidelity of individual plant species to specific habitats. These values will be calculated according to the procedure in the Manual using the list of plant species identified on the AA.

#### Calculation of Final Key Wildlife Habitat Ecological Integrity Assessment Score (Section 5)

The major components of the EIA include four core factors: landscape, soil/substrate, hydrology, and KWH and vegetation composition. The previously scored metrics that pertain to these core factors should be entered into the Scoring Form. Use these values to calculate the Overall KWH Ecological Integrity Assessment Score using the scale on the Scoring Form.

**Use the check boxes on the Scoring Form to note if any of the additional features are present** from the sources indicated as described in the Manual (Sections 3.5 and 5.1). **If the EIA score is not "Excellent"**, add additional points for unique resources present at the project area according to the instructions on the Scoring Form to calculate the Final Key Wildlife Habitat Ecological Integrity Assessment Score and Rating for the AA.